JOURNAL OF THE MARINE BIOLOGICAL ASSOCIATION

CONTENTS: VOLUME 71

Number 1, pp. 1-246, February 1991

PAGE

Comparative locomotor activity patterns in the portunid crabs *Liocarcinus holsatus* and *L. depurator*. Abelló, P., Reid, D.G. and Naylor, E. 1
Distribution patterns and dispersal of prosobranch gastropods along a seamount chain in the Atlantic Ocean. Leal, J.H. and Bouchet, P. 11
Behaviour, irrigation and respiration in *Eudistylia vancouveri* (Polychaeta: Sabellidae). Giangrande, A. 27
Virus-like particles in bacteria symbiotic in bivalve gills. Southward, E.C. and Southward, A.J. 37
The butterfly squid *Stoloteuthis leucoptera* in the Mediterranean. Orsi Relini, L. and Massi, D. 47
Larval metamorphosis of the opisthobranch mollusc *Adalaria proxima* (Gastropoda: Nudibranchia): the effects of choline and elevated potassium ion concentration. Todd, C.D., Bentley, M.G. and Havenhand, J.N. 53
Respiration and biometry in the sea cucumber *Holothuria forskali*. Astall, C.M. and Jones, M.B. 73
Fast pressure pulses and communication between fish. Gray, J.A.B. and Denton, E.J. 83
The development of bass, *Dicentrarchus labrax*, eggs in relation to temperature. Jennings, S. and Pawson, M.G. 107
Eggs of the *Microchirus* species (Soleidae) from the Bay of Algiers. Marinaro, J.Y. 117
Distribution and movements of intertidal butterfish *Pholis gunnellus*. Koop, J.H. and Gibson, R.N. 127
The ultrastructure of the flagellar root system of *Isochrysis galbana* (Prymnesiophyta). Hori, T. and Green, J.C. 137
Replicability of an epibenthic sampler. Brattegard, T. and Fosså, J.H. 153
Environmental Change ~ Biological Response: papers from meeting 167
Tolerances to high temperature of infaunal bivalves and the effect of geographical distribution, position on the shore and season. Wilson, J.G. and Elkaim, B. 169
Comparison of long-term trends in benthic and pelagic communities of the North Sea. Austen, M.C., Buchanan, J.B., Hunt, H.G., Josefson, A.B. and Kendall, M.A. 179
Variations in time and space: is biogeography relevant to studies of long-time scale change? Angel, M.V. 191
Relationship between a drop of sardine and albacore catches and a relaxation of wind, tested by a superposed epoch analysis. Littaye Mariette, A. 207
A comparison of some methods for analysing changes in benthic community structure. Warwick, R.M. and Clarke, K.R. 225
Book Notices 245

Number 2, pp. 247-494, May 1991

PAGE

Mechanisms of resistance to polychlorinated biphenyls (PCB) in two species of marine diatoms. Cohen, M.K., West, A.S., Cosper, E.M. and Wurster, C.F. 247

The *Ampelisca diadema* group of species (Amphipoda: Gammaridea) in British and Irish waters. Myers, A.A. and McGrath, D. 265

The Institute of Oceanographic Sciences Biology Programme in the Porcupine Seabight: background and general introduction. Rice, A.L., Billett, D.S.M., Thurston, M.H. and Lampitt, R.S. 281

Aspects of the biology of the deep-sea crab *Geryon trispinosus* from the Porcupine Seabight. Atrill, M.J., Hartnoll, R.G. and Rice, A.L. 311

Deep demersal fish assemblage structure in the Porcupine Seabight (eastern North Atlantic): slope sampling by three different trawls compared. Merrett, N.R., Gordon, J.D.M., Stehmann, M. and Haedrich, R.L. 329

Deep demersal fish assemblage structure in the Porcupine Seabight (eastern North Atlantic): results of single warp trawling at lower slope to abyssal soundings. Merrett, N.R., Haedrich, R.L., Gordon, J.D.M. and Stehmann, M. 359

Population biology of *Liocarcinus depurator* (Brachyura: Portunidae) in mussel raft culture areas in the Ría de Arousa (Galicia, NW Spain). Fernández, L., González-Gurriarán, E. and Freire, J. 375

The occurrence of *Ophelia bicornis* (Polychaeta) in and near the estuary of the River Exe, Devon. Harris, T. 391

Seasonal and spatial variation of epilithic microalgal distribution and abundance and its ingestion by *Patella vulgata* on a moderately exposed rocky shore. Hill, A.S. and Hawkins, S.J. 403

Development, growth, and survival of *Lepeophtheirus salmonis* (Copepoda: Caligidae) under laboratory conditions. Johnson, S.C. and Albright, L.J. 425

Carbon and nitrogen scope for growth as a function of diet in the sea scallop *Placopecten magellanicus*. Grant, J. and Cranford, P.J. 437

Trophic sources and pathways to the developing gametes of *Pecten maximus* (Bivalvia: Pectinidae). Le Pennec, M., Beninger, P.G., Dorange, G. and Paulet, Y.-M. 451

Image analysis of faecal material grazed upon by three species of copepods: evidence for coprorhexy, coprophagy and coprochaly. Noji, T.T., Estep, K.W., MacIntyre, F. and Norrbin, F. 465

The effect of synchronous moulting on body copper and zinc concentrations in four species of talitrid amphipods (Crustacea). Weeks, J.M. and Moore, P.G. 481

Short Communication:

Testing the validity of the synthesis ratio of protein to low molecular weight metabolites as an estimation of phytoplankton growth in the field. Madariaga, I. de, Fernández, E. and Serret, P. 489

Book Notices 493

Number 3, pp. 495-748, August 1991

PAGE

Forty years of changes in species composition and population density of barnacles on a rocky shore near Plymouth. Southward, A.J. 495
Factorial analysis in the ecology of rocky subtidal areas near Brest (West Brittany, France). Castric-Fey, A. and Chassé, C. 515
Foraging activity of limpets in normal and abnormal tidal regimes. Little, C., Partridge, J.C. and Teagle, L. 537
Biological and physical factors controlling the spatial distribution of the intertidal alga *Gelidium pristoides* in the eastern Cape, South Africa. Carter, A.R. and Anderson, R.J. 555
Effects of a toxic bloom of *Chrysochromulina polylepis* on the common dog-whelk, *Nucella lapillus*, on the Swedish west coast. Robertson, A. 569
Seasonal changes in epipelic diatoms along an intertidal shore, Berrow Flats, Somerset. Oppenheim, D.R. 579
A reassessment of the genus *Strombidinopsis* (Ciliophora, Choreotrichida) with descriptions of four new planktonic species and remarks on its taxonomy and phylogeny. Lynn, D.H., Montagnes, D.J.S., Dale, T., Gilron, G.L. and Strom, S.L. 597
Two new mesopelagic appendicularians: *Inopinata inflata* gen. nov., sp. nov., *Mesopelagica caudaornata* gen. nov., sp. nov. Fenaux, R. and Youngbluth, M.J. 613
The hypobranchial gland in *Morula granulata* (Gastropoda: Prosobranchia). Srilakshmi, G. ... 623
Effects on dietary lipids of the marine bivalve *Scrobicularia plana* feeding in different modes. Bradshaw, S.A., O'Hara, S.C.M., Corner, E.D.S and Eglinton, G. 635
Sexually dimorphic free-swimming behaviour in the amphipod crustacean *Ampelisca abdita*. Borowsky, B. and Aitken-Ander, P. 655
The influence of photoperiodicity on hatching of *Sepia officinalis*. Paulij, W.P., Herman, P.M.J., Roozen, M.E.F. and Denucé, J.M. 665
Size fluctuation in the eggs and newly hatched larvae of captive turbot (*Scophthalmus maximus*). McEvoy, L.-A. and McEvoy, J. 679
Fatty acid composition of early non-feeding larvae of the European flat oyster, *Ostrea edulis*. Helm, M.M., Holland, D.L., Utting, S.D. and East, J. 691
Abstracts: Mechanics and physiology of animal swimming. Rayner, J.M.V. 707
Short Communication:
Satellite tracking of a loggerhead turtle (*Caretta caretta*) in the Mediterranean. Hays, G.C., Webb, P.I., Hayes, J.P., Priede, I.G. and French, J. 743
Book Notices 747

Number 4, pp. 749-991, November 1991

PAGE

Colour polymorphism in relation to spatial distribution in some intertidal isopods in northern Spain. Arrontes, J. 749
A motographic analysis of foraging behaviour in intertidal chitons (*Acanthopleura* spp.) Santini, G., Della Santina, P. and Chelazzi, G. 759
Some aspects of the specific habitat requirements of *Ophelia bicornis* (Polychaeta). Harris, T. 771
Influence of temperature on embryonic and larval development in *Necora puber* (Brachyura, Portunidae). Valdés, L., Alvarez-Ossorio, M.T. and González-Gurriarán, E. 787
The age structure and growth dynamics of young-of-the-year bass, *Dicentrarchus labrax*, populations. Jennings, S., Lancaster, J.E., Ryland, J.S. and Shackley, S.E. 799
The uptake and accumulation of zinc and copper from solution by two species of talitrid amphipods (Crustacea). Weeks, J.M. and Rainbow, P.S. 811
The American oyster drill, *Urosalpinx cinerea* (Gastropoda): evidence of decline in an imposex-affected population (R. Blackwater, Essex). Gibbs, P.E., Spencer, B.E. and Pascoe, P.L. 827
Plant-animal interactions in the marine benthos. Williams, G.A. and Hawkins, S.J. 839
The logic of ecological experiments: a case history from studies of the distribution of macro-algae on rocky intertidal shores. Underwood, A.J. 841
Refugial *Laminaria* abundance and reduction in urchin grazing in communities in the north-west Atlantic. Keats, D.W. 867
Effect of encrusting bryozoans on the physiology of their algal substratum. Muñoz, J., Cancino, J.M. and Molina, M.X. 877
Seasonal variation in *Pilayella littoralis* (Phaeophyceae) and its consequences as a food source for the amphipod, *Gammarus lawrencianus*, in the intertidal of Newfoundland. Steele, D.H. and Whittick, A. 883
Amphipod food preference and *Iridaea* spp. (Rhodophyta) spore release and dispersal. Buschmann, A.H. 891
Interactions between the amphipod *Corophium volutator* and macroalgal mats on estuarine mudflats. Raffaelli, D., Limia, J., Hull, S. and Pont, S. 899
Short Communication:
Recent studies on the fecundity of the European flat oyster *(Ostrea edulis)* spawning stock in the Solent. Utting, S.D., Helm, M.M. and Millican, P.F. 909
Announcement 913
Book Notices 915
Annual Report 917
Financial Statement 985

CORRIGENDUM, VOLUME 70

Page 807, Eqn(1) *for* $\ln(F) = -6{\cdot}7 + 3{\cdot}20 \ln(L)$ *read* $\ln(F) = -6{\cdot}07 + 3{\cdot}20 \ln(L)$